CAS No.	3159-62-4	1592-23-0	64755-01-7	557-04-0	Mixture <sup>14</sup>	4499-91-6	4485-12-5	53422-16-5	7620-77-1	68783-36-8	64754-95-6	38900-29-7	68603-11-2	
Name:		Octadecanoic	Fatty acids,	Supporting	Supporting	Docosanoic	Octadecanoic	Octadecanoic	Octadecanoic	Fatty acids,	Castor oil,	Nonanedioic	Hydrocarbon	
	acid, 12-hydroxy-	acid calcium	tallow, calcium	Chemical:	Chemical	acid, lithium sal	t acid, lithium salt	acid, 12-hydroxy		$C_{16-22}$ , lithium		acid, dilithium	waxes,	
	calcium salt (2:1)	salt	salts	Octadecanoic				, methyl ester,	, monolithium	salts	lithium salt	salt	petroleum,	1
				acid				lithium salt	salt				oxidized, Me	1
				magnesium salt	t								esters, calcium	Read Across Range to
													salts	Untested Category Member
Synonym:	Calcium 12-	Calcium		Magnesium	Generic	Lithium	Lithium stearate	Lithium 12-	Lithium 12-			Dilithium		
	hydroxystearate	stearate		stearate	calcium	docosanoate		hydroxystearate	hydroxystearate			azelate		1
					complex grease	•		(same as 7620-	(same as 53422	<del>'- </del>				1
								77-1)	16-5)					
Endpoint		·	T	Т	PHYSICAL-C	HEMICAL PROF		Т		T	T	1		T
Melting Point (°C)	320	179				271	249		264			186		179 to 320
Boiling Point (°C)	730	661				624	578		611			484		484 to 730
Vapor Pressure (hPa) Partition Coefficient	1 x 10E-21 11.7	6 X 10E-14 14.3				5 x 10E-14 6.1	1 x 10E-12 4.1		2 x 10E-16 2.6			2 x 10E-9 -3.6		<1 x 10E-5 -3.6 to 14.3
Water Solubility <sup>1</sup> (mg/L)	9.7 x 10E-9	40 at 15 C				0.04	4.1	222 (0.1	_			(877)		<0.001 to 877
water Solubility (mg/L)	(6.4 x 10E-7)					(2.0 x 10E-5)	(0.002)	(0.1)	<b>′</b>			(011)		
		<u> </u>	<u> </u>	<u> </u>	1	FNV	IRONMENTAL F	ATF		<u> </u>	<u> </u>	<u> </u>		
Photodegradation, OH <sup>-</sup> reaction														0.2 to 1.4
$\Gamma_{1/2}$ (d)	0.2	0.2				0.4	0.5		0.4			1.4		
Stability in Water	stable	stable				stable	stable		stable			stable		stable
Environ. Distribution	<0.1% air	<0.1% air				<0.1% air	<0.1% air		<0.1% air			<0.1% air		air <0.1%
	<0.1% water	<0.1% water				<0.1% water	8% water		73% water			>99% water		soil <0.1% to 98% water <0.1% to 99%
	98% soil	98% soil				98% soil	90% soil		26% soil			<0.1% soil		water <0.1% to 99% sediment <0.1% to 2%
Piodogradation algorification	2% sediment	2% sediment				2% sediment	2% sediment		0.6% sediment			<0.1% sediment		Sediment <0.170 to 270
Biodegradation classification	inherently	readily					readily							<u> </u>
						ENVIR	ONMENTAL EFI	FECTS						
Acute Fish, mg/L		no toxicity at				no toxicity at						no toxicity at		
	LC50 >12,500	limits of solubility				limits of solubility	,	LC50 = 123	LC50 > 2000 mg/L	_		limits of solubility		LC50 >2000 mg/L <sup>2</sup>
Exposure type	) Whole grosse	Ecosar Model				Ecosar Model		Ecosar Model	thickener OWD			Ecosar Model		
	whole grease	thickener				thickener		thickener	thickener, OWD			thickener		1
Acute Invertebrate, mg/L		no toxicity at				no toxicity at						no toxicity at		
	EL50 >1000	limits of solubility				limits of solubility	,					limits of solubility		EL50 >1000 mg/L <sup>3</sup>
Exposure type	1	Faccar Madel				Ecosar Model						Ecosar Model		
	WAF, whole grease	thickener				thickener						thickener		
Algae, mg/L	>1000	no toxicity at				no toxicity at						no toxicity at		$EL_b 50 = >100 \text{ to } >1000 \text{ mg/L}^3$
	$EL_r 50 = >320 \text{ to}$	limits of solubility				limits of solubility	,					limits of solubility		$EL_r 50 = >320 \text{ to } >1000 \text{ mg/L}$
Exposure type	)	Ecosar Model				Ecosar Model						Ecosar Model		
. ,	WAF, whole grease	thickener				thickener						thickener		1
							1							
A	1	ľ	1	L L D FO 40 000	T	Н	EALTH EFFECT	I		1	1	Т	<u>r</u>	Т
Acute (mg/kg, oral)				LD50 >10,000				LD50 >5,000	LD50 >5,000			LD50 >5,000		LD50 >5,000
Acute (mg/kg, dermal)								LD50 >3,000 <sub>6</sub>	LD50 >3,000 <sub>6</sub>			LD50 >3,000 <sub>6</sub>		LD50 >2,000
Repeated-Dose (mg/kg/day, oral)				NOEL = 5% of										
Deposited Dags (mar/land)				diet (~2,500)				NOAEL = 1,000	NOAEL = 1,000			NOAEL = 1,000		NOAEL = 1,000
Repeated-Dose (mg/kg/day, dermal)								NOAEL = 2,100	NOAEL = 2,100	0		7		1
demai)								NOEL = 2,100	NOEL = 2,100			8		1
					4.			$NOEL = 2,000^{1}$	$NOEL = 2,000^{1}$	NOEL=2,000 <sup>1</sup>		NOAEL = 2,100		1
					NOAEL = $2,000^{14}$	+		NOAEL <500 <sup>12</sup>	NOAEL <500 <sup>12</sup>	NOAEL <500 <sup>12</sup>		NOEL = 2,100		NOAEL = 2,000
Genotoxicity, in vitro Genotoxicity, in vivo				negative			1		1	1	1			negative
Reproductive toxicity (dermal,	1		1	1					+	1	1			negative
mg/kg)														NOAEL = $2,000^{13}$
Developmental toxicity (dermal,														
mg/kg)	1		1	1				NOEL = 2,000 <sup>11</sup>	NOEL = 2,000 <sup>11</sup>	NOEL = 2,000 <sup>11</sup>				NOAEL = 2,000
Carcinogenicity (dermal, mg,								negative (50	negative (50					negative
mouse skin painting)	<u> </u>		ļ	1			1	mg/dose)	mg/dose)	1	1	1		<b></b>
	<u> </u>					lank colle indicate	no data Value fe	om the road seres	s rango will be use	d for those sells				
	Footnotes:				В	ialik celis indicate	e no uata. Value fr	om me read across	s range will be used	u for these cells.				
		EPΔ 2000\ provide	e two actimates fo	r water colubility	alues. The first value	ie nrecente the set	imate using the rela	ationship with Kou /	Meylan and Howard	1994) while the v	alue in naronthoa	se wae calculated b	w the fragment con	stant method (Meylan and Howard
	1995).	∟⊢∧, ∠uuu) piovide	รอ เพบ <del>ย</del> อแทเสเยร ใ0	ı water surubility Va	aiucs. IIIE IIIS(Väll	ie hreserirs riie esti	imate using the fela	mononih mini vom (I	ivieyian anu noward,	, 1994) Wille (ne V	aiue iii paieiimese	so was calculated D	y une magment cons	stant method (weylan and noward
	,	od ac an ail water	dispersion (OMD)											
	Whole grease test Invertebrate and a		. ,		testing of a whole	arease								
	5 NOAEL is for a gre					•								
							90 davs in rats with	sample R96000257	75.					
								sample R96000257						
								h sample R9600025						
	Test was with PAF	RL-3093-GR-81, a	grease containing 7	7.5% lithium 12-hyd	droxystearate.	-								
							3 weeks in rats with							
	•					•		eric Lithium Grease					00/	100
	1	_	se that contained 7	.9% CAS No. 7620	9-77-1 and 0.9% C	45 No. 68783-36-8	. Study was in male	e ⊢344 rats with "Bo	rated-Lithium 12-Hyd	droxy Complex Ge	eneric Grease". G	rease contained ~2	U% performance ac	dditives, over twice the amount
	typically used in gre	eases.												

CAS No.	3159-62-4	1592-23-0	64755-01-7	557-04-0	Mixture <sup>14</sup>	4499-91-6	4485-12-5	53422-16-5	7620-77-1	68783-36-8	64754-95-6	38900-29-7	68603-11-2	
Name:	Octadecanoic	Octadecanoic	Fatty acids,	Supporting	Supporting	Docosanoic	Octadecanoic	Octadecanoic	Octadecanoic	Fatty acids,	Castor oil,	Nonanedioic	Hydrocarbon	
	acid, 12-hydroxy-,	acid calcium	tallow, calcium	Chemical:	Chemical	acid, lithium salt	acid, lithium salt	acid, 12-hydroxy-	acid, 12-hydroxy-	C <sub>16-22</sub> , lithium	hydrogenated,	acid, dilithium	waxes,	
	calcium salt (2:1)	salt	salts	Octadecanoic				, methyl ester,	, monolithium	salts	lithium salt	salt	petroleum,	
				acid				lithium salt	salt				oxidized, Me	
				magnesium salt									esters, calcium	Read Across Range to
													salts	Untested Category Members
Synonym:	Calcium 12-	Calcium		Magnesium	Generic	Lithium	Lithium stearate	Lithium 12-	Lithium 12-			Dilithium		
	hydroxystearate	stearate		stearate	calcium	docosanoate		hydroxystearate	hydroxystearate			azelate		
					complex grease			(same as 7620-	(same as 53422-					
								77-1)	16-5)					
	13 Results from repeat dose studies and developmental studies with lithium complex greases with typical additive concentrations can be used to read across to all members of category.													
	A 13-weeks dermal toxicity study was performed on a "generic calcium complex grease" that contained 3.5% CAS No. 62-54-4 (acetic acid, calcium salt), 3.5% CAS No. 64754-97-8 (fatty acids, coco, calcium salts), 1.4% CAS No. 69012-90-4 (fatty acids, C6-12, calcium salts),													
	and 1.2% CAS No. 66071-81-6 (fatty acids, tallow, hydrogenated, calcium salts). The structures of the thickeners in this grease were judged to be similar enough to those in the Lubricating Grease Thickeners category that the data from this study can be applied by read-across.													